

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



ACC029-11

会場:102

時間:5月26日 15:45-16:00

## 氷床コアの水同位体比を用いた「南極ドームふじ」と「その水蒸気起源」温度の復元

Reconstructions of temperature histories of the Dome Fuji site and its moisture source using water isotope records

植村 立<sup>1\*</sup>, Valerie Masson-Delmotte<sup>2</sup>, Jean Jouzel<sup>2</sup>, Amaelle Landais<sup>2</sup>, Francoise Vimeux<sup>2</sup>, 本山 秀明<sup>3</sup>, Barbara Stenni<sup>4</sup>  
Ryu Uemura<sup>1\*</sup>, Valerie Masson-Delmotte<sup>2</sup>, Jean Jouzel<sup>2</sup>, Amaelle Landais<sup>2</sup>, Francoise Vimeux<sup>2</sup>, Hideaki Motoyama<sup>3</sup>, Barbara Stenni<sup>4</sup>

<sup>1</sup> 琉球大学 理学部 海洋自然科学科, <sup>2</sup>LSCE, CEA/CNRS/UVSQ, <sup>3</sup>情報・システム研究機構 国立極地研究所, <sup>4</sup>University of Trieste

<sup>1</sup>University of the Ryukyus, <sup>2</sup>LSCE, CEA/CNRS/UVSQ, <sup>3</sup>NIPR, <sup>4</sup>University of Trieste

A stable isotope ratio (D/H or  $^{18}\text{O}/^{16}\text{O}$ ) of the polar ice cores is widely used as an air-temperature proxy. Further, a combined use of these isotopes provides a parameter, deuterium-excess ( $d$ ), and provides the information on the ocean surface conditions in the moisture source for polar precipitation. Here we re-evaluate several coefficients used for reconstructing Antarctic site temperature and temperature in the moisture source regions. The new coefficients were applied to the revised D/H and  $d$  records of Dome Fuji core which cover past 360 kyr period. Differences between this study and previous estimations will be discussed.