

## Acknowledgments

The author would like to express his sincere gratitude to Prof. Toshifumi Mukai for his continuous guidance to write this paper. The author also wishes to express his sincere thanks to Prof. Masaki Ejiri for his continuous guidance and helps for the author to participate in various important projects, the AKEBONO satellite, the 30th Japanese Antarctic Research Expedition (JARE-30), and so on. Without his help, the author could not imagine any ideas about the subjects treated in this paper. The author also wishes to express many thanks to all the members of the AKEBONO satellite project, especially to Prof. Koichiro Tsuruda, the project manager, Prof. Hiroshi Oya, the project scientist, Prof. Takasi Oguti, the PI of ATV team, Dr. Eisuke Kaneda, and Dr. Tatsundo Yamamoto. He also wishes to express his hearty thanks to all the members of JARE-30, especially to Dr. Noriyuki Kurihara and Mr. Akira-Sessai Yukimatu. The author is deeply grateful to all the stuffs of the upper atmosphere physics group in the National Institute of Polar Research (NIPR), especially to Prof. Takeo Hirasawa for his continuous encouragement, Prof. Natuo Sato, Prof. Hisao Yamagishi, and Prof. Takehiko Aso. Special thanks are due to Dr. Masakazu Watanabe for fruitful discussions, Prof. Takayuki Ono for giving many instructions about the system in the Auroral Data Center in NIPR to the author, all the co-workers in the Data Center, and Prof. Yohsuke Kamide and Prof. Ryoichi Fujii for their encouragement. The author also would like to express his special gratitude to all the members of the upper atmospheric science division in British Antarctic Survey (BAS), especially to Dr. Mike Pinnock, Dr. Alan Rodger, and Dr. John Dudeney. Without the precious stay at BAS in Cambridge, his scientific motivation should be very different.

The AKEBONO ATV-UV data were provided from the database of the Institute of Space and Astronautical Science (ISAS). The AKEBONO LEP, EFD, and MGF science database data were provided by Prof. Toshifumi Mukai, Dr. Ayako Matsuoka, Prof. Hajime Hayakawa and Prof. Koichiro Tsuruda, and Dr. Tsutomu Nagatsuma and Prof. Hiroshi Fukunishi. The NOAA satellite data, DMSP satellite SSJ/4 data and GOES satellite data were supplied from the NGDC, and the IMP-8 data were supplied from the NSSDC. The high time resolution GOES satellite magnetic field data were provided by Prof. Tsugunobu Nagai, and the DMSP satellite SSIES data were provided by Dr. Marc Hairston. The LANL satellite energetic particle data were provided by Dr. Geoff Reeves. All-sky camera film data were provided by Dr. F. T. Berkey for the South Pole data, by Dr.

A. Y. Feldstein for the Vostok data, and by Dr. M. Pinnock for the Halley data. As for South Pole station, the photometer and riometer data were provided by Dr. T. J. Rosenberg, and the magnetometer data were provided by Dr. L.J. Lanzerotti, respectively. The Halley HF-radar data were provided by Dr. M. Pinnock. The Halley magnetometer and riometer data and the Faraday magnetometer data were provided by Dr. Matt Palley. The Sanae magnetometer and riometer data were provided by Dr. P. R. Sutcliffe and Dr. Mike Mathews, respectively. The Maitri magnetometer data were provided by Mr. Ajay Dhar. The Magnetometer data at Mawson, Davis and Casey were provided by Dr. G. Burns, and the induction magnetometer data at Hermanus were provided by Dr. P. R. Sutcliffe. The author thanks Dr I.R. Mann and Dr D.K. Milling for the SAMNET data. SAMNET is a PPARC National Facility deployed and operated by the University of York. He also thanks Dr. T. Moretto for the magnetometer data from the Greenland stations which were maintained by Danish Meteorological Institute. The author also used the magnetometer data from the EISCAT magnetometer cross stations which were maintained by the Technical University of Braunschweig, and the magnetometer data from the Canadian stations which were maintained by the Geological Survey of Canada and provided by Mr. Gerrit Jansen van Beek. Other magnetometer data from all over the world were provided by Prof. T. Iyemori from the WDC for geomagnetism, Kyoto. The observations at Syowa and Asuka stations were carried out by JARE-30 leading by Prof. M. Ejiri. The auroral observations were carried out by the wintering member including the author at Syowa station and by Mr. A.-S. Yukimatu at Asuka station.